

# CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# Slasher\* ORGANIC\* WEEDKILLER

**ACTIVE CONSTITUENT: 525 g/L NONANOIC ACID** 


For non-selective control of seedling and young broadleaf and grass weeds, for suppression of established weeds and perennial species and for control of moss and algae as per the directions for use.

- ✓ Versatile weed, moss and algae killer
- $\diagup$  Visible results within 1 hour on most weeds $^{\dagger}$   $\checkmark$
- √ Glyphosate free
- \* Registered for use in organic farming systems

contents 20 L

ORGANIC CROP PROTECTANTS

1956 Dandenong Rd, Clayton VIC 3168 Telephone: 1800 634 204 www.ocp.com.au



**Active made from natural plant oils** 







Not approved under USDA-NOP Certification



### DIRECTIONS FOR USE

### RESTRAINTS:

DO NOT spray if weeds are wet.

DO NOT apply if rainfall is expected before spray dries on plant surfaces.

DO NOT water treated area for 24 hours to achieve maximum results.

DO NOT apply if heavy rains or storms that are likely to cause runoff from soil are forecast within 3 days of application.

DO NOT irrigate to the point of runoff from soil for at least 3 days after application.

DO NOT allow spray to contact or drift onto plants you do not want killed.

FOR PROFESSIONAL USE ONLY.

SITUATION	WEEDS	RATE	CRITICAL COMMENTS
OUTDOOR AREAS Paths, Driveways, Gardens, Around sheds, Amenity horticulture areas, Protected cropping situations, Around Nursery stock. Spot Spraying in Lawns and Turf.	Seedling and young annual and perennial weeds and grasses including fat hen, potato weed, stinking roger, couch grass, chickweed, lambs tongue, peppercress, capeweed, common storksbill, clover. More advanced or established weeds may only be suppressed and require repeated application.	70 mL per litre of water.  Mix 7 litres per 100 litres water. Apply 900-1400 L spray per ha.	Spray to completely cover weed surface. Reapplication at around 7 days may be necessary if regrowth occurs. This product may damage lawn and turf when used as a spot spray. Perennial weed species may require repeated applications to obtain long term control. To control weeds in Protected cropping situations apply on each side of the crop row taking care not to allow spray to contact any of the crop foliage. Apply as a band spray 400 mm wide on each side of the order of which was a band spray 400 mm wide on each side of the order of which was a band spray 400 mm wide on each side of the order of which was a band spray 400 mm wide on each side of the order of which was a band spray 400 mm wide on each side of the order of which was a band spray 400 mm wide on each side of the order of which was a band spray 400 mm wide on each side of the order of which was a band spray 400 mm wide on each side of the order of which was a band spray 400 mm wide on each side of the order of which was a band spray 400 mm wide on each side of the order of which was a band spray 400 mm wide on each side of the order of which was a band spray 400 mm wide on each side of the order of which was a band spray 400 mm wide on each side of the order o
GENERAL USES Including around buildings, Commercial and Industrial areas, Domestic and Public Service areas. Rights-of way. Bushland & Forests.			
AGRICULTURAL AREAS			
Orchards and Vineyards.			
Fallow soil.			
Gardens, Paths, Rockeries, Tiled Roofs, Walls, Driveways, Around buildings, Spot Spraying in Lawns and Turf.	Algae, Lichen, Liverwort, Moss.	70 mL per litre of water.	Spray to completely cover plant surface. For driveways or heavy infestations, clean off by brushing after 3 days and reapply if necessary.

## NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

### **GENERAL INSTRUCTIONS:**

Slasher® Organic\* Weedkiller is a natural contact weedkiller containing pelargonic acid (nonanoic acid). Pelargonic acid occurs naturally in plants including geraniums. It acts on the foliage, causing it to quickly dehydrate and collapse. Results are seen as browning and withing usually within 1 hour of application.

¹ Visible results can take longer than 1 hour on certain tough weeds or weeds in shaded conditions. For best results, apply when temperature is over 10°C. To be effective, Slasher® Organic\* Weedkiller must come into contact with the green parts of the plant. Slasher® Organic\* Weedkiller will control many weed species and is non-selective.

MIXING: Half fill spray tank and commence agitation. Slowly add required amount of Salsher® Organic\* Weedkiller. Top up with required quantity of water. Agitate well before commencing spray procedure. Do not mix more than is needed.

APPLICATION: Ensure that all weed foliage is totally covered with spray as Slasher® Organic\* Weedkiller is a contact spray. Partial coverage will give only partial control.

Apply by ground equipment only using for example boom or spot sprayers. Apply using a coarse droplet spray. Optimum application volumes are related to weed maturity and density. Contact with the weeds is essential for control, as such apply in sufficient water to ensure complete and thorough coverage of foliage.

DO NOT spray porous surfaces without prior testing.

May cause discolouration on porous hard surfaces.

PRECAUTIONS: Wear eye protection when mixing or using.

RE-ENTRY PERIOD: Do not enter treated areas until product has dried.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS: DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT: DO NOT contaminate wetlands or watercourses with this product or used containers. Not compatible with integrated pest management (IPM) programs utilising beneficial arthropods. Minimise spray drift to reduce harmful effects on beneficial arthropods in non-crop areas.

STORAGE AND DISPOSAL: Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undituted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS: Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist for equivalent clothing, elbow-length chemical resistant gloves and face shield or goggles. When using the prepared spray wear gauntlet length chemical resistant gloves and goggles. Wash hands after use. After each day's use wash gloves, face shield or goggles, and contaminated clothing.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. If swallowed, on NOT induce vomiting. Give a glass of water. If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

# FOR EMERGENCIES ONLY CALL: 1800 033 111 (24 HOURS).

SDS: Additional information is listed in the Safety Data Sheet available from the supplier and occ.com.au.

May cause an allergic skin reaction

Slasher and the OCP drop device are trade marks of Organic Crop Protectants.

APVMA Approval No. 68118/121803