

TriCure^{AD}

Soil Surfactant

CONTENTS

U.S. Volume		U.S. Weight		Metric Volume		Metric Weight	
<input type="checkbox"/>	1.0 Gal. (3.7 L)	8.7 Lbs.	(3.9 kg)	<input type="checkbox"/>	5.0 L	5.2 kg	
<input type="checkbox"/>	2.5 Gal. (9.4 L)	21.8 Lbs.	(9.8 kg)	<input type="checkbox"/>	10.0 L	10.4 kg	
<input type="checkbox"/>	5.0 Gal. (18.9 L)	43.6 Lbs.	(19.7 kg)	<input type="checkbox"/>	115.0 L	120.1 kg	
<input type="checkbox"/>	30.0 Gal. (113.5 L)	261.6 Lbs.	(118.6 kg)	<input type="checkbox"/>	200.0 L	209.0 kg	
<input type="checkbox"/>	52.9 Gal. (200.2 L)	461.2 Lbs.	(209.2 kg)	<input type="checkbox"/>	1000.0 L	1041.3 kg	
<input type="checkbox"/>	260.0 Gal. (984.2 L)	2267.2 Lbs.	(1028.3 kg)				

Density: 8.69 lbs/gal at 68°F

CONTAINS NON-PLANT-FOOD INGREDIENTS

Active Ingredient:

100% Oxirane 2-methyl polymer with oxirane



User Guide enclosed. SDS available at tricure.com

KEEP OUT OF REACH OF CHILDREN



WARNING

Irritant to eyes, skin, and ingestion.

Use proper gloves and goggles when handling and wash hands before touching eyes or eating. Get medical advice/attention if you feel unwell. Dispose of contents/container in accordance with local/regional/national and international regulations.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.html>

221216

Mitchell Products LLC

1205 W. Main Street
Millville, NJ 08332

856.327.2005 | tricure.com

MITCHELL
PRODUCTS

DIRECTIONS FOR USE

For best results, **TriCure AD**® applications should begin in early spring, before first signs of heat-related stress appear. **TriCure AD** can also be applied curatively on existing dry areas and LDS. In warmer climates, application should be year-round or when environmental stress conditions begin to impact turf performance. Immediate watering-in is not necessary to prevent burning as **TriCure AD** is non-phytotoxic. For best results, heavy irrigation is recommended when applying to facilitate movement of chemical into the soil.

Spray Application

Applications can be made monthly or on a seasonal basis (effective approximately three months). Refer to the Spray Application Rates section to determine the appropriate rate. (Note: To maintain an even, optimum surfactant level in the soil, monthly applications are preferable.)

Application by Injection

Application of **TriCure AD** via irrigation lines is recommended where possible as surfactant is optimally disbursed and reduced application rates with comparable results are also often possible. Recommended rates are 5lt per Hectare.

TANK MIXING

Although **TriCure AD** can be mixed with most chemicals used in turf management, it is recommended that field-testing be done on a small area before spraying to assure safety concerns are met.

PRECAUTIONARY STATEMENTS

- Overexposure causes eye irritation, excessive watering, redness.
- Overexposure to skin may cause itching and redness.
- Harmful if swallowed. May cause diarrhea, nausea and vomiting.
- Wear approved eye protection and clothing.
- Keep out of reach of children.

IN CASE OF CONTACT

- **EYES** If eye exposure occurs, wash with water for at least 15 minutes. See a physician.
- **SKIN** Eye and skin irritant. Wash skin and clothing thoroughly with water after exposure.
- **INGESTION** Contact physician or poison-control center. If victim is conscious and able to swallow, quickly give milk or water to dilute. Do not give sodium bicarbonate, vinegar, or fruit juices. Never give anything by mouth if victim is unconscious or having convulsions. Induce vomiting only upon advice of a physician.

SPRAY APPLICATION RATES

ml/100m ² 10 lt. Water	Usage
120 ml / Monthly	Regular use for normal conditions
180 ml / Monthly	High stress conditions and curative for LDS
First application should be in early spring, or when first signs of heat stress appear.	